

Herz 2015 · [Suppl 2] 40:182–189
 DOI 10.1007/s00059-014-4185-z
 Received: 18 August 2014
 Revised: 13 October 2014
 Accepted: 2 November 2014
 Published online: 3 December 2014
 © Urban & Vogel 2014

E. Şahan · M. Gül · S. Şahan · E. Sokmen · Y.A. Guray · O. Tufekçioğlu

Department of Cardiology, Atatürk Pulmonary Disease and Thorax Surgery Research Hospital, Ankara

Pseudoaneurysm of the mitral–aortic intervalvular fibrosa

A new comprehensive review

Abbreviations	
LVOT	Left ventricular outflow tract
MAIVF	Mitral–aortic intervalvular fibrosa
P-MAIVF	Pseudoaneurysm of the mitral–aortic intervalvular fibrosa
TEE	Transesophageal echocardiography
TTE	Transthoracic echocardiography

The mitral–aortic intervalvular fibrosa (MAIVF) is an avascular structure located adjacent to the left ventricular outflow tract between the aortic and mitral valve providing functional and anatomical integrity.

Pseudoaneurysm of the MAIVF (P-MAIVF) is a rare but potentially life-threatening condition. With the increasing incidence of infective endocarditis and cardiac surgery, a likely increment in the frequency of pseudoaneurysm is expected. Transesophageal echocardiography (TEE) has facilitated the visualization of this area and enhanced the detection and assessment of pseudoaneurysm.

P-MAIVF is often associated with infection or surgical trauma. A congenital origin and blunt chest trauma are among the other causes. While it is likely to follow an asymptomatic course, symptoms of endocarditis, heart failure, chest pain, shortness of breath, and cerebrovascular events are common clinical presentations.

The recommended treatment is surgery. However, conservative therapy is an alternative approach for high-risk patients

or those who refuse surgery with a reasonable follow-up period with echocardiography of 6–12 months.

Methods

We found 487 articles in the PubMed database by searching the keywords “pseudoaneurysm of the mitral–aortic intervalvular fibrosa,” “left ventricular outflow tract aneurysm,” “left ventricular pseudoaneurysm,” and “subvalvular aneurysm”. All of the articles were screened manually and 101 were selected due to their relevance to P-MAIVF. These articles were published between 1960 and March 2014, and one was in French. A total of 161 cases were evaluated in 104 articles. The cases were classified according to their age, sex, etiology, clinical presentation, diagnosis, complication, treatment, and survival status. Besides these 161 patients collected in our literature review, our hospital established a definite diagnosis in another series of eight patients (■ Tab. 1). Since only five of them remained unpublished, we included them in our comprehensive review, thereby reaching a total of 166 cases.

Etiology and pathophysiology

The MAIVF is bounded by the left and right fibrous trigones, thus gaining a functional and anatomical integrity between the aortic valve and mitral valve. The roof is formed by the pericardium, while its base is situated adjacent to the

left ventricular outflow tract (LVOT). Half of the noncoronary cusp, some part of the left coronary cusp, and the anterior mitral leaflet constitute an immediate vicinity of this area.

Since the MAIVF is relatively avascular, it becomes more prone to infections and traumas. Infection of this area may result in abscess or pseudoaneurysm formations. The opening of the abscess formation into the LVOT creates a suitable milieu for the formation of P-MAIVF.

The most common causative factors are infective endocarditis and surgical trauma. Aortic valve infections may extend directly into the MAIVF area. The relatively avascular nature of the MAIVF renders it more prone to infectious insults and hence abscess formations. Should the abscess sac communicate with the LVOT, a pseudoaneurysm can possibly be generated.

Aortic regurgitation is another entity that contributes to the formation of pseudoaneurysm. Infected aortic regurgitation jet impinging on this innately sensitive region may also predispose to the development of pseudoaneurysm. The combination of bicuspid aortic valve and PMAIVF seems a common condition because of the congenital compromise in this area (bicuspid/native valve ratio: 0.44 in our review).

Additional material online: This article includes an additional Table. You will find this supplemental at dx.doi.org/10.1007/s00059-014-4185-z.

Tab. 1 Clinical and imaging features of our patients												
Case	Gender	Age (years)	Time from diagnosis to surgery	Size	Diagnostic method	Aortic valve	Endocarditis	Causative microorganism	Comorbidity	Treatment	Surgical details	
1	Female	35	No surgery	2.1×1.1 cm	TTE/3DTTE	Native	–	–	Takayasu	Refused surgery; lost to follow-up	–	
2	Female	23	No surgery	1.2×3.1 cm	TTE/3DTTE	Native	Aortic valve endocarditis	<i>S. aureus</i>	–	Exitus	–	
3	Male	43	38 months	3.2×1.9 cm	MDCT/TTE/TEE	AVR	–	–	–	Surgery	AVR + closure of pseudoaneurysm with synthetic graft	
4	Female	42	26 months	1.6×2.3 cm	TTE/TEE	AVR	–	–	–	Surgery	Closure of pseudoaneurysm with synthetic graft	
5	Male	68	108 months	0.9×2.1 cm	TTE/TEE	AVR	–	–	MVR	Surgery	Simple closure of pseudoaneurysm neck	
6	Male	71	No surgery	2.7×2.1 cm	TTE/TEE/3D	Native	Aortic valve endocarditis	<i>S. aureus</i>	–	Refused surgery; noncardiac exitus	–	
7	Male	55	14 months	2.8×1.6 cm	TTE/TEE	Bioprosthes	–	–	Severe MR	Surgery	AVR + MVR + closure of pseudoaneurysm with pericardial graft	
8	Male	72	76 months	1.1×1.4 cm	MDCT/TTE/TEE	AVR	Metallic aortic valve endocarditis	<i>S. aureus</i>	–	Surgery	AVR + simple closure of pseudoaneurysm neck	

TTE transthoracic echocardiography, TEE transesophageal echocardiography, MDCT multidetector computed tomography, AVR aortic valve replacement, MVR mitral valve replacement

In the selected articles, 68 patients had active endocarditis (40%), whereas 30 patients had a history of previous endocarditis (18%). The causative microorganisms were reported to be *Staphylococcus* species in 22 patients [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 102, 104], *Streptococcus* species in 14 patients [1, 7, 17, 18, 19, 20, 21, 22, 23, 103], *Bacillus* species in one patient [24], *Brucella suis* in two patients [25, 26], *Escherichia coli* in one patient [27], and *Enterococcus* species in four patients [17, 28, 29, 30]. Fungal infections were the associated etiology in two patients, in one of whom it was *Paecilomyces lilacinus* [31] and in the other, *Monilia albicans* [32].

MAIVF can become subject to damage due to valvular surgery. Not only valvular surgeries, but other surgical and interventional procedures such as cardiac catheterization, radiofrequency ablation, and ventricular septal defect repair in the vicinity of this area can also result in P-MAIVF. In our literature review we found that P-MAIVF developed in 87 patients (52%) following aortic surgery, among whom 52 also had had a history of active or prior infective endocarditis.

Congenital heart diseases are another causative factor of P-MAIVF. Cardiovascular defects such as ventricular septal defect, atrial septal defect, patent ductus arteriosus, aortic coarctation, and bicuspid

aortic valve may be detected at the time of diagnosis [47, 73].

Clinical presentation

P-MAIVF often remains asymptomatic in the absence of complications. It is usually diagnosed during a routine examination. The most commonly encountered clinical situations are symptoms related to an active infection, shortness of breath, and symptoms/signs of heart failure [16, 32, 33]. Chest pain, cerebrovascular events, and systemic embolism are other likely clinical presentations [16, 32, 33]. The most common clinical presentation in the reviewed articles was active en-

docarditis, with 68 patients in this clinical situation [2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 14, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 45, 46, 47, 102, 103, 104], 36 patients had been admitted with shortness of breath and/or other heart failure-related symptoms/signs [6, 27, 29, 30, 31, 32, 34, 41, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60]. Five patients had presented with cerebrovascular events [4, 7, 10, 21, 46], and 19 patients had described typical or atypical angina [9, 16, 20, 58, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72]; 17 patients had been asymptomatic at the time of diagnosis [1, 49, 60, 61, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83]. In 33 patients the reports did not reveal adequate information regarding the symptomatology. A chest wall mass gained attention as an unusual clinical presentation in one patient [84].

Complications

Impingement by the pseudoaneurysm of the surrounding tissues leads to a number of complications. Specifically, progressive enlargement of the pseudoaneurysm may compress the coronary arteries, thus resulting in angina or even myocardial infarction [12, 21, 62, 64, 65, 67, 72, 79, 85]. The most common compression is on the left circumflex coronary artery. Other coronary arteries can also be affected. Compression of the pulmonary artery [47] may result in pulmonary hypertension. Compression of the mitral annulus and anterior mitral valve leaflet is likely to result in mitral regurgitation [51].

Fistula formations into adjacent cardiac structures constitute another complication. A fistula may develop between the LVOT and the left atrium or the aorta. Upon formation of a fistula between the LVOT and the left atrium, an eccentric flow can be visualized mimicking a mitral regurgitant jet. Accordingly, any new eccentric mitral regurgitation detected in addition to a previous or recent aortic valve replacement should raise suspicion of fistula formation.

If the pseudoaneurysm ruptures into the pericardium, fatal complications may ensue [78, 102]. The most typical example of this situation is pericardial tamponade. Purulent and hemorrhagic pericarditis

Herz 2015 · [Suppl 2] 40:182–189 DOI 10.1007/s00059-014-4185-z
© Urban & Vogel 2014

E. Şahan · M. Gül · S. Şahan · E. Sokmen · Y.A. Guray · O. Tufekçioğlu Pseudoaneurysm of the mitral–aortic intervalvular fibrosa. A new comprehensive review

Abstract

Pseudoaneurysm of the mitral–aortic intervalvular fibrosa (P-MAIVF) is an infrequent but potentially life-threatening condition. Both transthoracic echocardiography (TTE) and transesophageal echocardiography (TEE) can detect P-MAIVF with sensitivity rates of 43 and 90%, respectively. The typical finding of echocardiography is a pulsatile echo-free sac that expands in systole and collapses in diastole. Our review comprises 166 patients with P-MAIVF, including eight cases in our hospital and 158 cases from the literature. P-MAIVF is often associated with infection or surgical trauma. While it is likely to maintain an asymptomatic course, symptoms of shortness of breath, heart failure, val-

vular disease, chest pain, endocarditis, and cerebrovascular events are common clinical presentations. The recommended treatment is surgery. However, conservative therapy is an alternative approach for high-risk patients or when surgical treatment is refused. With the increasing incidence of cardiac surgery and infective endocarditis, a likely increment in the new diagnosis of pseudoaneurysm is expected.

Keywords

Pseudoaneurysm · Mitral–aortic intervalvular fibrosa · Transthoracic echocardiography · Transesophageal echocardiography · Therapy

Pseudoaneurysma der mitral-aortalen intervalvulären Fibrosa. Eine neue umfassende Übersicht

Zusammenfassung

Das Pseudoaneurysma der mitral-aortalen intervalvulären Fibrosa (MAIVF, aortomitraler Übergang) ist nicht häufig, aber potenziell lebensbedrohlich. Sowohl mit der transthorakalen Echokardiographie (TTE) als auch mit der transösophagealen Echokardiographie (TEE) kann ein Pseudoaneurysma der MAIVF (P-MAIVF) mit einer Sensitivitätsrate von 43 bzw. 90% diagnostiziert werden. Typischer Befund bei der Echokardiographie ist ein pulsierender echofreier Sack, der sich in der Systole ausdehnt und in der Diastole zusammenfällt. In die Studie wurden 166 Patienten mit P-MAIVF, darunter 8 Fälle aus der Klinik der Autoren und 158 Fälle aus der Literatur, aufgenommen. Oft geht ein P-MAIVF mit einer Infektion oder einem chirurgischen Trauma einher. Es ist zwar wahrscheinlich, dass der Verlauf asymptomatisch ist, aber Symp-

tome wie Luftnot, Herzinsuffizienz, Herzklap-penerkrankungen, Schmerzen in der Brust, Endokarditis und zerebrovaskuläre Ereignisse sind häufige klinische Zeichen. Zumeist wird die chirurgische Behandlung empfohlen. Die konservative Therapie ist jedoch ein alternativer Ansatz für Hochrisikopatienten oder in Fällen, in denen die Operation abgelehnt wird. Mit Zunahme der Häufigkeit von Herzoperationen und infektiöser Endokarditis ist auch von einem Anstieg der neuen Diagnose des Pseudoaneurysmas auszugehen.

Schlüsselwörter

Pseudoaneurysma · Mitral-aortale intervalvuläre Fibrosa · Transthorakale Echokardiographie · Transösophageale Echokardiographie · Therapie

is another complication of P-MAIVF. In one case report, P-MAIVF was reported to have ruptured into the anterior chest wall, generating a pulsatile mass in the chest wall [84].

Clots may occur within the cavity of the pseudoaneurysm, and they may be associated with cerebrovascular events and systemic emboli.

Diagnosis

Echocardiography is the most widely used method in the diagnosis of the pseudoaneurysm. Transthoracic echocardiography (TTE) was performed in 152 of 166 cases (92%) (■ Fig. 1, 2). Using TTE, a definitive diagnosis could be made in 60 patients (36%); suspicious lesions were observed in 24 patients; and P-MAIVF was not evident in 68 patients. Moreover, transesophageal echocardiography




Fig. 1 ▲ Four-chamber view of a pseudoaneurysm (asterisk)




Fig. 2 ▲ Parasternal long-axis view of a pseudoaneurysm (asterisk)




Fig. 3 ▲ Transesophageal echocardiographic view of a pseudoaneurysm (arrow)




Fig. 4 ▲ Color Doppler view of a pseudoaneurysm (arrow)

(TEE) was also implemented for a further detailed examination in most of the patients (66%) undergoing TTE. P-MAIVF was evident in all patients who were evaluated using TEE. Even if a large pseudoaneurysm could be depicted in TTE, TEE was necessary for the definitive diagnosis and detection of the complications. TEE proved superior to TTE in identifying the pseudoaneurysm (■ Fig. 3). One study demonstrated that the diagnostic sensitivity of TTE was markedly lower than that of TEE (43% vs. %90, $p < 0.05$) [46].

The typical finding of echocardiography is a pulsatile echo-free sac that expands in systole and collapses in diastole. This almost always plays a major role in the definitive diagnosis of P-MAIVF [6].

Color Doppler imaging can help establish the diagnosis on the basis of forward blood flow into the pseudoaneurysm during systole and backflow into the LVOT

during diastole (■ Fig. 4). In addition, color Doppler imaging helps demonstrate the relationship between the pseudoaneurysm and other heart structures [6]. Color Doppler examination relies on the turbulent flow inside the cavity and can prove useful in revealing any communication between the echo-free space, namely, the pseudoaneurysm, and the LVOT. The source and direction of the blood flow can be defined more clearly using color Doppler TEE imaging, which further facilitates the diagnosis of any complications.

Furthermore, compared with two-dimensional echocardiography, three-dimensional echocardiography provides a better anatomic and morphological visualization of the cardiac structures (■ Fig. 5). Three-dimensional echocardiography is likely to be more useful, especially if two-dimensional views are insufficient for identifying the patho-anatom-

ic structures in recognizing the complications of endocarditis [35, 74].

Other diagnostic methods are computed tomography, magnetic resonance imaging, and left ventricular catheterization. Computed tomography was used for the purpose of a definitive diagnosis in 24 patients (■ Fig. 6, [9, 12, 15, 29, 36, 37, 48, 59, 61, 65, 66, 69, 71, 72, 73, 78, 80, 82, 87, 88]); magnetic resonance imaging in 11 patients [2, 37, 51, 62, 77, 81, 83, 89, 90]; and, the diagnosis was confirmed by left heart catheterization in 19 patients [12, 21, 32, 37, 46, 47, 52, 64, 67, 69, 70, 76, 85, 89, 91, 92, 103].

Differential diagnosis

Aortic root abscess should be kept in mind in the differential diagnosis. The characteristic echocardiographic feature of P-MAIVF, namely, the pulsatile echo-free pouch, is not evident in aortic root



Fig. 5 ◀ Three-dimensional view of a pseudoaneurysm (asterisk)

abscess. Moreover, Color Doppler imaging demonstrates no inflow to the aortic root abscess. Furthermore, such kinds of abscesses are prone to be smaller than P-MAIVF.

Characteristic ante- and retrograde blood flow visualized using color Doppler imaging P-MAIVF may also be confused with aortic stenosis and aortic regurgitation.

When the pseudoaneurysm communicates with the left atrium, its flow can be confused with mitral regurgitation jet. Accordingly, suspicions regarding P-MAIVF should rise whenever an eccentric jet has been visualized within the left atrium, especially in patients with the history of a previous infective endocarditis or aortic valve replacement surgery. Furthermore, the origin of such a jet is likely to be elucidated more thoroughly using TEE thanks to its better spatial and temporal resolution.

Management and treatment

Surgery

The clinical outcome of P-MAIVF is hard to predict. Rupture of the pseudoaneurysm into the pericardium may be fatal and hence when the pseudoaneurysm is diagnosed, surgical treatment should be recommended to all patients even if they are asymptomatic. A number of surgical procedures have been used, including: simple closure of the aneurysm neck; closure using pericardial or synthetic graft materials; and complete removal of the pseudoaneurysm sac from the LVOT. In

most of the cases, aortic valve replacement or aortic root surgery were included in the surgical repair of P-MAIVF.

Successful percutaneous closure attempts were preferred in some cases, with uneventful follow-ups reported [27, 65, 77].

Type of surgery

Most of the reviewed patients underwent surgical treatment. Pseudoaneurysm repair encompassed concomitant aortic surgery including aortic valve replacement and/or aortic root replacement in 71 patients [2, 6, 7, 8, 9, 12, 13, 14, 15, 19, 20, 21, 22, 23, 24, 26, 30, 31, 32, 38, 41, 42, 43, 45, 46, 48, 49, 51, 52, 53, 62, 63, 64, 67, 68, 88, 92, 93, 94, 95, 104]; concomitant aortic valve replacement and mitral valve replacement in 11 patients [2, 5, 7, 14, 18, 33, 50, 54, 55, 56]; and concomitant mitral valve replacement in only one patient [29]. On the other hand, pseudoaneurysm repair was implemented without any concomitant valve surgery in 23 patients [3, 16, 17, 33, 37, 47, 59, 69, 70, 72, 76, 82, 83, 84, 87, 91, 96, 97, 98, 103]; coronary arterial by-pass grafting was added to the repair procedure in two patients [64, 67]. One patient had a successful orthotopic cardiac transplantation [67].

Follow-up without surgery

An alternative approach to surgical correction is conservative therapy. Surgical correction might be high risk for various reasons. In particular, comorbidities could cause contraindications for surgery, and these patients should be followed up with



Fig. 6 ▲ Computed tomography view of a pseudoaneurysm (asterisk)

medical treatment. The natural course of P-MAIVF remains to be clearly elucidated. For this reason, close clinical and echocardiographic follow-up is essential. The combination of TTE and TEE should be recommended in all follow-ups. Pseudoaneurysm rupture has been reported only rarely, but one must be vigilant in terms of complications. Surgical treatment was not used in 29 patients [1, 4, 17, 29, 33, 36, 39, 46, 47, 49, 60, 61, 71, 73, 74, 78, 80, 81, 99]. The follow-up periods for these patients ranged from 10 months to 9 years, with a mean of 30 months. The dimensions of the pseudoaneurysm did not change in ten patients. On the other hand, the size of the pseudoaneurysm increased progressively in some patients. In one patient with a 9-year follow-up, the size of the pseudoaneurysm increased from 0.6 cm to 4.9×7.8 cm [1]. Interestingly, this patient remained asymptomatic.

Mortality

Mortality occurred in 29 patients during follow-up, which was due to: heart failure in five patients [34, 58]; coronary arterial disease or coronary artery compression-related complications in three patients; [1, 21, 79]; P-MAIVF rupture into the pericardium in three patients [33, 44, 102]; infection and/or sepsis in five patients [2, 4, 17, 28, 40]; cerebrovascular accident in one patient [35]; and intraoperatively or postoperatively in 11 patients [2, 14, 17, 21, 30, 41, 47, 59, 60, 64]; in one case autopsy was refused by the patient's family [60].

Sixteen patients had active infective endocarditis or a history of endocarditis;

of these patients, eight had aortic valve surgery, five were diagnosed at autopsy, and one had congenital heart disease. The cause of death in two patients could not be found in the articles.

Discussion and conclusion

As mentioned, P-MAIVF is a rare clinical entity. Postmortem reports regarding some African patients were the first documents to which we ever gained access in the literature, wherein the exact etiology had remained unknown [58]. Thanks to the increasing recognition and usage of echocardiography modalities around the world, the number of diagnoses has been rising progressively so that increasing insights into the etiology are being gained.

The most common causes are infection and surgical trauma. Both native and prosthetic valve endocarditis may result in P-MAIVF. In some cases the pseudoaneurysm develops due to the opening of an abscess formation through the LVOT. P-MAIVF was also suggested to be associated with insult incurred by the aortic regurgitation jet trauma toward the MAIVF area.

The most common traumatic surgery is aortic valve replacement. Since MAIVF is closely knitted with the aortic root anatomy, it may be easily damaged during such surgeries. Therefore, any eccentric mitral regurgitation identified in a patient with previous or recent aortic valve replacement should be further evaluated for a probable P-MAIVF. Aortic valve replacement is not the only reason of surgical trauma. The same may hold true following a mitral valve replacement surgery. Additionally, ventricular septal defect repair, intracardiac radiofrequency catheter ablation, and other cardiac structural repair operations are also likely to result in P-MAIVF during follow-up. Such patients, therefore, should be regarded as high risk and be approached in the relevant manner.

Surgical repair is the treatment of choice, and the most frequently preferred surgical method comprises pseudoaneurysm repair together with aortic surgery. Simple closure, repair using a graft, and percutaneous repair are among the alternative treatment methods. A conservative

approach is preferred when surgery is not considered. Patients should be monitored regularly in reasonable time lines, namely, 6–12 months, using a combination of TTE and TEE.

The increasing incidence of infective endocarditis and aortic surgeries in recent years is expected to expand the current series of P-MAIVF cases. Therefore, clinicians should always bear in mind the likelihood of P-MAIVF developing in such cases.

Study limitations

Our study was limited by some incomplete or and/or missing patient reports and the relatively small sample size. Furthermore, we only reviewed the English- and Turkish-language literature and one French article regarding this rare disease entity.

Corresponding address

E. Şahan

Department of Cardiology, Atatürk Pulmonary Disease and Thorax Surgery Research Hospital Sanatoryum Street, 06280 Ankara Turkey
ekremsahan@hotmail.com

Compliance with ethical guidelines

Conflict of interest. E. Şahan, M. Gül, S. Şahan, E. Sokmen, Y.A. Guray, and O. Tufekçioğlu state that there are no conflicts of interest.

The accompanying manuscript does not include studies on humans or animals.

References

1. Gin A, Hong H, Rosenblatt A et al (2011) Pseudoaneurysms of the mitral-aortic intervalvular fibrosa: survival without reoperation. *Am Heart J* 161(1):130 e1–e5
2. Kassim TA, Lowery RC, Nasur A et al (2010) Pseudoaneurysm of mitral-aortic intervalvular fibrosa: two case reports and review of literature. *Eur J Echocardiogr* 11(3):E7
3. Kunavarapu C, Olkovsky Y, Lafferty JC et al (2008) Unusual cardiac complications of *Staphylococcus aureus* endocarditis. *J Am Soc Echocardiogr* 21(2):187 e3–e5
4. Penugonda N, Duncan K, Afonso L (2007) Complex endocarditis in an immunocompromised host: the role of three-dimensional echocardiography. *J Am Soc Echocardiogr* 20(11):1314 e9–e11

5. Ashmeik K, Pai RG (2000) An unusual case of acquired ventricular septal defect as a complication of aortic valve endocarditis: echocardiographic delineation of multiple subvalvular complications in one patient. *J Am Soc Echocardiogr* 13(7):693–695
6. Cheong YM, Hamer AW, Johns JA (1997) Mitral-aortic intervalvular fibrosa pseudoaneurysm complicating aortic valve endocarditis. *Aust N Z J Med* 27(2):193–194
7. Bansal RC, Graham BM, Jutzy KR et al (1990) Left ventricular outflow tract to left atrial communication secondary to rupture of mitral-aortic intervalvular fibrosa in infective endocarditis: diagnosis by transesophageal echocardiography and color flow imaging. *J Am Coll Cardiol* 15(2):499–504
8. Sadat K, Joshi D, Sudhakar S et al (2012) Incremental role of three-dimensional transesophageal echocardiography in the assessment of mitral-aortic intervalvular fibrosa abscess. *Echocardiography* 29(6):742–744
9. Schaap J, Brinkman EB, Heijmen RH (2011) Left ventricular outflow tract pseudoaneurysm compromising blood flow through the left main coronary artery after mechanical aortic valve implantation. *Eur Heart J* 32(12):1508
10. Koch R, Kapoor A, Spencer KT (2003) Stroke in patient with an intervalvular fibrosa pseudoaneurysm and aortic pseudoaneurysm. *J Am Soc Echocardiogr* 16(8):894–896
11. Moaref A, Shahrzad S, Aslani A (2009) Acquired aortoventricular tunnel: a rare complication of infective endocarditis. *Echocardiography* 26(1):82–83
12. Berrizbeitia LD, Anderson WA (1997) Ultrafast computed tomography in infectious pseudoaneurysm of the left ventricular outflow tract. *J Thorac Cardiovasc Surg* 114(1):138–139
13. Lastowiecka E, Biederman A, Szajewski T, Rydlewska-Sadowska W (2002) The rupture of peri-aortic infective aneurysm into the left atrium and the left ventricular outflow tract: preoperative diagnosis by transthoracic echocardiography. *Echocardiography* 19(3):173–176
14. Leung DY, Cranney GB, Hopkins AP, Walsh WF (1994) Role of transoesophageal echocardiography in the diagnosis and management of aortic root abscess. *Br Heart J* 72(2):175–181
15. Malvindi PG, Ornaghi D, Tarelli G, Raffa GM (2012) Left ventricular pseudoaneurysm following aortic valve prosthesis endocarditis. *J Cardiovasc Med (Hagerstown)* 13(7):457–459
16. Sudhakar S, Sewani A, Agrawal M, Uretsky BF (2010) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa (MAIVF): a comprehensive review. *J Am Soc Echocardiogr* 23(10):1009–1018
17. Karalis DG, Bansal RC, Hauck AJ et al (1992) Transesophageal echocardiographic recognition of subaortic complications in aortic valve endocarditis. Clinical and surgical implications. *Circulation* 86(2):353–362
18. Dores H, Abecasis J, Ribeiros R et al (2012) Uncommon acquired Gerbode defect following extensive bicuspid aortic valve endocarditis. *Cardiovasc Ultrasound* 10:7
19. Takimoto K, Arai F, Kita T, Sasaki S (2010) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa on a native aortic valve following infective endocarditis. *J Anesth* 24(2):260–263
20. Tak T (2003) Pseudoaneurysm of mitral-aortic intervalvular fibrosa. *Clin Med Res* 1(1):49–52

21. Parashara DK, Jacobs LE, Kotler MN et al (1995) Angina caused by systolic compression of the left coronary artery as a result of pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Am Heart J* 129(2):417–421
22. Perelman MJ, Sugimoto J, Arcilla RA, Karp RB (1989) Aortic root replacement for complicated bacterial endocarditis in an infant. *J Pediatr Surg* 24(11):1121–1123
23. Frank MW, Mavroudis C, Backer CL, Rocchini AP (1998) Repair of mitral valve and subaortic mycotic aneurysm in a child with endocarditis. *Ann Thorac Surg* 65(6):1788–1790
24. Caselli S, Mazzezi G, Tritapepe L et al (2011) 3D echocardiographic delineation of mitral-aortic intervalvular fibrosa pseudoaneurysm caused by bicuspid aortic valve endocarditis. *Echocardiography* 28(1):E1–E4
25. Layman TE, January LE (1967) Mycotic left ventricular aneurysm involving the fibrous atrioventricular body. *Am J Cardiol* 20(3):423–427
26. Sharifkazemi MB, Moarref AR, Rezaian S, Rezaian GR (2013) Brucella endocarditis of pseudoaneurysm of an aortic composite graft. *J Cardiovasc Ultrasound* 21(4):183–185
27. Gorselen E van, Nihoyannopoulos P, Verhorst P et al (2011) Severe regurgitation due to perforation of the mitral-aortic intervalvular fibrosa 3 years after aortic valve replacement. *Eur J Echocardiogr* 12(4):E30
28. Araujo IR, Nunes Mdo C, Gelape CL et al (2012) Challenge in the management of infective endocarditis with multiple valvular involvement. *Rev Soc Bras Med Trop* 45(2):272–274
29. Grimaldi A, Ho SY, Pozzoli A et al (2011) Pseudoaneurysm of mitral-aortic intervalvular fibrosa. *Interact Cardiovasc Thorac Surg* 13(2):142–147
30. Velibey Y, Güngör B, Bolca O (2013) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa and complementary role of 3D transesophageal echocardiographic imaging. *Anadolu Kardiyol Derg* 13(8):E-44
31. Muller H, Cikirikcioglu M, Lerch R (2008) Subaortic aneurysm caused by *Paecilomyces lilacinus* endocarditis. *Arch Cardiovasc Dis* 101(11–12):803–804
32. Bansal RC, Moloney PM, Marsa RJ, Jacobson JG (1983) Echocardiographic features of a mycotic aneurysm of the left ventricular outflow tract caused by perforation of mitral-aortic intervalvular fibrosa. *Circulation* 67(4):930–934
33. Xie M, Li Y, Cheng TO et al (2013) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Int J Cardiol* 166(1):2–7
34. Panneerselvam A, Subbiahnaadar P (2011) Ruptured mitral-aortic intervalvular fibrosa aneurysm with flail aortic valve. *BMJ Case Rep*
35. Selcuk H, Selcuk MT, Tufekcioglu O, Arat N (2009) The use of three-dimensional echocardiography in the visualization of pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Turk Kardiyol Dern Ars* 37(5):328–331
36. Ghersin E, Litmanovich D, Agmon Y, Milo S (2005) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa following aortic valve replacement—diagnosis and dynamic evaluation with multidetector CT and transesophageal echocardiography. *Interact Cardiovasc Thorac Surg* 4(6):502–504
37. Yokoyama Y, Tamaki S, Kato N et al (2003) Pseudoaneurysm from the mitral-aortic intervalvular fibrosa following endocarditis. *Jpn J Thorac Cardiovasc Surg* 51(8):374–377
38. Griffiths BE, Petch MC, English TA (1982) Echocardiographic detection of subvalvar aortic root aneurysm extending to mitral valve annulus as complication of aortic valve endocarditis. *Br Heart J* 47(4):392–396
39. Pai RG, Ortega V, Ferry DR (1995) A case of mitral-aortic intervalvular fibrosa aneurysm with unique flow patterns and long-term natural survival. *J Am Soc Echocardiogr* 8(2):211–214
40. Pirat B, Buerger J, Zoghbi WA (2005) Echocardiographic features of pseudoaneurysm of the mitral-aortic inter-valvular fibrosa. *Anadolu Kardiyol Derg* 5(4):344–345
41. Takamiya Y, Miura S, Sako H et al (2007) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa following infective endocarditis in a patient with acute heart failure: a case report. *J Cardiol* 49(6):353–356
42. Garneau S, Demers P, Denault AY (2007) Unexpected periaortic cavity during heart surgery. *Anesth Analg* 104(1):75–76
43. Hori D, Tanaka M, Yamaguchi A et al (2008) Extracardiac aneurysm of the interleaflet triangle above the aortic-mitral curtain due to infective endocarditis of the bicuspid aortic valve. *Gen Thorac Cardiovasc Surg* 56(8):424–426
44. Saint-Martin P, Rogers C, Carpenter E et al (2009) Subaortic pseudoaneurysm of the left ventricle complicating staphylococcal endocarditis. *J Forensic Sci* 54(4):930–932
45. Sogabe O, Ohya T (2007) A case of tuberculous endocarditis with acute aortic valve insufficiency and annular subvalvular left ventricular aneurysm. *Gen Thorac Cardiovasc Surg* 55(2):61–64
46. Afridi I, Apostolidou MA, Saad RM, Zoghbi WA (1995) Pseudoaneurysms of the mitral-aortic intervalvular fibrosa: dynamic characterization using transesophageal echocardiographic and Doppler techniques. *J Am Coll Cardiol* 25(1):137–145
47. Gelehrter S, Wright G, Gless T et al (2002) Left ventricular outflow tract pseudoaneurysms in congenital heart disease. *Am J Cardiol* 90(7):806–809
48. Malcles G, Azarnoush K, Ravan R et al (2011) Pseudoaneurysm of mitro-aortic continuity: a rare complication after aortic valve replacement. *Ann Cardiol Angeiol (Paris)* 60(5):296–299
49. Kahveci G, Keles N, Bayrak F, Mutlu B (2008) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa after aortic valve replacement. *Turk Kardiyol Dern Ars* 36(7):499
50. Ivens EL, Thompson CR, Moss RR et al (2008) Images in cardiovascular medicine: prosthetic aortic valve and conduit dehiscence with large periconduit cavity, ascending aortic aneurysm and severe mitral regurgitation. *Eur J Echocardiogr* 9(1):148–151
51. Espinosa-Caliani JS, Montijano A, Melero JM, Montiel A (2000) Pseudoaneurysm in the mitral-aortic intervalvular fibrosa. A cause of mitral regurgitation. *Eur J Cardiothorac Surg* 17(6):757–759
52. Tiwari KK, Murzi M, Mariani M, Glauber M (2009) Giant pseudoaneurysm of the left ventricle outflow tract after aortic root replacement for extensive endocarditis. *Eur J Cardiothorac Surg* 36(2):399
53. Turiel M, Gottardi B, Muzzupappa S (2001) A case of subannular aortic aneurysm detected by transesophageal echocardiography. *Acta Cardiol* 56(6):395–397
54. Aikatt S, Lundergan CF, Adkins MS, Lewis JF (2003) Delayed presentation of left ventricular outflow tract aneurysm after penetrating cardiac trauma. *J Am Soc Echocardiogr* 16(10):1085–1087
55. Kruthoff WA, Akl BF, Blacky AR (1995) Surgical repair of an anterior mitral subvalvular aneurysm. *Ann Thorac Surg* 59(4):1001–1003
56. Harish A, Hatibu S, Ramkumar SR et al (2009) Aneurysm of mitral aortic intervalvular fibrosa—a case report and review. *Indian Heart J* 61(4):394–396
57. Moorthy N, Kumar S, Tewari S et al (2012) Mitral-aortic intervalvular fibrosa aneurysm with rupture into left atrium: an uncommon cause of acute dyspnea. *Heart Views* 13(1):13–15
58. Lurie AO (1960) Left ventricular aneurysm in the African. *Br Heart J* 22:181–188
59. Acioli Pereira L, Fontes Gontijo P, Alcantara Farran J et al (2013) Giant pseudoaneurysm of the left ventricular outflow tract: a rare disease. *Rev Port Cardiol* 32(6):541–544
60. Parakh N, Yadav N, Kathuria S et al (2013) Pseudoaneurysm of mitral aortic intervalvular fibrosa: two case reports. *Indian Heart J* 65(1):100–103
61. Hasin T, Reisner SA, Agmon Y (2011) Large pseudoaneurysms of the mitral-aortic intervalvular fibrosa: long-term natural history without surgery in two patients. *Eur J Echocardiogr* 12(3):E24
62. Kim HW, Chung CH (2010) Mitral-aortic intervalvular fibrosa pseudoaneurysm resulting in the displacement of the left main coronary artery after aortic valve replacement. *J Thorac Cardiovasc Surg* 139(2):e18–e20
63. Rodrigues Borges AG, Suresh K, Mirza H et al (2002) False aneurysm of the mitral-aortic intervalvular fibrosa after uncomplicated aortic valve replacement. *J Am Soc Echocardiogr* 15(7):743–745
64. Antonellis J, Kostopoulos K, Routoulas T et al (1997) Aneurysm of the mitral-aortic intervalvular fibrosa as a rare cause of angina pectoris: angiographic demonstration. *Cathet Cardiovasc Diagn* 42(4):423–426
65. Romaguera R, Slack MC, Waksman R et al (2010) IMAGE RADIO MED: Percutaneous closure of a left ventricular outflow tract pseudoaneurysm causing extrinsic left coronary artery compression by transseptal approach. *Circulation* 121(4):e20–e22
66. Turkvatan A, Demirkan B, Guray Y (2011) Multidetector computed tomographic angiography diagnosis of a giant pseudoaneurysm of the left ventricular outflow tract. *Eur J Cardiothorac Surg* 39(3):e30
67. Almeida J, Pinho P, Torres JP et al (2002) Pseudoaneurysm of the mitral-aortic fibrosa: myocardial ischemia secondary to left coronary compression. *J Am Soc Echocardiogr* 15(1):96–98
68. Aoyagi S, Fukunaga S, Otsuka H et al (2004) Left ventricular outflow tract pseudoaneurysm after aortic valve replacement: case report. *J Heart Valve Dis* 13(1):145–148
69. Linhartova K, Veselka J, Adla T (2007) Left ventricular pseudoaneurysm as a late complication of mitral annuloplasty. *Eur Heart J* 28(19):2360
70. Jaggers J, Fullerton DA, Campbell DN, Clarke DR (1996) LVOT pseudoaneurysm: a late complication of extended aortic root replacement. *Ann Thorac Surg* 62(4):1198–1199
71. Tsai IC, Hsieh SR, Chern MS et al (2009) Pseudoaneurysm in the left ventricular outflow tract after prosthetic aortic valve implantation: evaluation upon multidetector-row computed tomography. *Tex Heart Inst J* 36(5):428–432

72. Joshi SS, Jagadeesh AM, Furtado A, Bhat S (2013) Transesophageal echocardiography in surgical management of pseudoaneurysm of mitral-aortic intervalvular fibrosa with aneurysms of right sinus of Valsalva and left main coronary artery. *Ann Card Anaesth* 16(1):40–43
73. Tsai IC, Fu YC, Lin PC et al (2009) MDCT evaluation of congenital mitral-aortic intervalvular fibrosa aneurysm: implications for the aetiology and differential diagnosis. *Pediatr Radiol* 39(1):80–83
74. Tufekcioglu O, Ozlu MF, Cay S et al (2008) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa in a patient with Takayasu's arteritis. *Can J Cardiol* 24(9):718
75. Gupta R, Jammula P, Huang MH et al (2006) An unusual complication after aortic valve replacement. *J Clin Ultrasound* 34(7):361–364
76. Yamamoto S, Tedoriya T, Masuda Z et al (2006) Mitral-aortic intervalvular pseudoaneurysm with ventricular septal defect. *Asian Cardiovasc Thorac Ann* 14(2):e35–e37
77. Jimenez Valero S, Garcia E, Gonzalez Pinto A, Delcan JL (2005) Percutaneous closure of pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Rev Esp Cardiol* 58(12):1473–1475
78. Gungor B, Altay S, Dayi SU, Bolca O (2012) 3D echocardiographic evaluation of ruptured pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Anadolu Kardiyol Derg* 12(8):E42–E43
79. Normann SJ (1991) Annular subaortic aneurysm resulting in sudden death. *Clin Cardiol* 14(1):68–72
80. Dogan M, Vural MG, Turkvatan A, Yeter E (2012) Incidentally diagnosed pseudoaneurysm of mitral-aortic intervalvular fibrosa. *Anadolu Kardiyol Derg* 12(8):E41–E42
81. Barranhas AD, Dias MC, Dos Santos AA et al (2012) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa presenting after chest trauma and diagnosed by cardiac magnetic resonance: a case report. *J Med Case Rep* 6(1):357
82. Spampinato RA, Borger MA, Strottdrees E, Mohr FW (2013) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa as a complication after minimally invasive mitral valve repair. *Interact Cardiovasc Thorac Surg* 16(3):396–398
83. Choh NA, Shaheen F, Rather H, Aslam K (2013) Pseudoaneurysm of mitral-aortic intervalvular fibrosa in a child: Demonstration by MDCT and MRI. *Ann Pediatr Cardiol* 6(1):80–82
84. Agirbasli M, Fadel BM (1999) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa: a long-term complication of infective endocarditis. *Echocardiography* 16(3):253–257
85. Matthews RV, French WJ, Criley JM (1989) Chest trauma and subvalvular left ventricular aneurysms. *Chest* 95(2):474–476
86. Sousa L, Branco L, Abreu J et al (1999) A pseudoaneurysm of the mitral-aortic intervalvular fibrosis after aortic valve replacement. *Rev Port Cardiol* 18(2):177–178
87. Da Col U, Ramoni E, Di Bella I, Ragni T (2009) An unusual left ventricular outflow pseudoaneurysm: usefulness of echocardiography and multidetector computed tomography for surgical repair. *Cardiovasc Intervent Radiol* 32(1):188–191
88. Entrikin DW, Shroff GS, Kon ND, Carr JJ (2011) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa: a delayed complication of aortic root replacement. *J Cardiovasc Comput Tomogr* 5(5):333–335
89. Winkler ML, Higgins CB (1986) MRI of perivalvular infectious pseudoaneurysms. *AJR Am J Roentgenol* 147(2):253–256
90. Robles P, Sonlleve A (2007) Aneurysm of the intervalvular mitroaortic fibrosa after aortic valve replacement diagnosed by cardiovascular magnetic resonance imaging. *Intern Med* 46(21):1825
91. Bolcal C, Baysan O, Doganci S et al (2007) A mitral-aortic intervalvular fibrosa false aneurysm in a patient without overt heart disease. *Can J Cardiol* 23(5):389–390
92. Goldfarb B, Rovner M, Goldbraich N (1997) Pseudoaneurysm of the left ventricular outflow tract with reentry into the ascending aorta: an iatrogenic left ventricular ascending aortic fistula. *J Am Soc Echocardiogr* 10(8):866–868
93. Ueno T, Sakata R, Iguro Y, Yamamoto H (2008) Intervalvular fibrosa pseudoaneurysm with projectile shunt flow to left atrium. *Interact Cardiovasc Thorac Surg* 7(4):727–729
94. Strauch JT, Wippermann J, Krep H, Wahlers T (2008) Subannular perforation after long-lasting aortic valve replacement mimicking mitral insufficiency. *Ann Thorac Surg* 85(1):311–313
95. Economopoulos G (2007) Delayed pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Ann Thorac Surg* 83(6):2233
96. Han J, He Y, Li Z et al (2009) Pseudoaneurysm of the mitral-aortic intervalvular fibrosa in a patient after radio frequency catheter ablation of atrial fibrillation. *J Ultrasound Med* 28(2):249–251
97. Pongratz G, Pohlmann M, Gehling G, Bachmann K (1997) Images in cardiovascular medicine. Pseudoaneurysm in the intervalvular mitral-aortic region after endocarditis and prosthetic aortic valve replacement. *Circulation* 96(9):3241–3242
98. Dawson D, Pessotto R, Neary P (2009) Inter-valvular fibrosa pseudo-aneurysm as a late complication after aortic valve surgery. *Eur J Echocardiogr* 10(1):169–170
99. Salerno D, Donati G, Forconi S, Gori T (2008) Giant pseudoaneurysm of the mitro-aortic intervalvular fibrosa: incidental diagnosis. *Intern Emerg Med* 3(3):279–282
100. Frary G, Hasselman T, Patel P (2012) Atypical left ventricular outflow tract aneurysm diagnosed by three-dimensional echocardiography. *Cardiol Young* 22(4):478–480
101. Kahveci G, Yilmaz F, Mutlu B (2008) Bicuspid aortic valve endocarditis complicated with aortic valve aneurysm and pseudoaneurysm of the mitral-aortic intervalvular fibrosa. *Echocardiography* 25(7):799–801
102. Ekici F, Kocabas A, Aktas D et al (2014) Native aortic valve endocarditis complicated by pseudoaneurysm of mitral-aortic intervalvular fibrosa. *Echocardiography* 31(2):60–63
103. Silbiger JJ, Krasner A, Chikwe J et al (2013) Pseudoaneurysm formation in infective endocarditis. *Echocardiography* 30(10):319–321
104. Chandra S, Ameta D, Kharwar RB et al (2013) Three-dimensional echocardiographic delineation of an acquired aorto-left atrial fistula complicating native aortic valve endocarditis—“advantage of three dimensions”. *Echocardiography* 30(10):326–330