

On The New Hyperbolic Wave Solutions to Wu-Zhang System Models

Tuğba YAZGAN

Ataturk University: Ataturk Universitesi

Esin ILHAN

Ahi Evran Universitesi Muhendislik Mimarlik Fakultesi

Ercan ÇELİK (✉ ercelik@atauni.edu.tr)

Ataturk Universitesi <https://orcid.org/0000-0002-1402-1457>

Hasan BULUT

Firat Üniversitesi: Firat Universitesi

Research Article

Keywords: The Sine-Gordon expansion method (SGEM), The modified expansion function method (MEFM), Wu-Zhang system

Posted Date: February 9th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1184408/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Optical and Quantum Electronics on April 17th, 2022. See the published version at <https://doi.org/10.1007/s11082-022-03683-y>.

On The New hyperbolic Wave Solutions to Wu-Zhang System Models

Tuğba YAZGAN^a, Esin ILHAN^b, Ercan ÇELİK^{a,1}, Hasan BULUT^c

^a Ataturk University, Faculty of Science, Department of Mathematics, Erzurum-TURKEY

^b Kırşehir Ahi Evran University, Faculty of Engineering and Architecture, Kırşehir, Turkey

^c Fırat University, Faculty of Science, Department of Mathematics, Elazığ-TURKEY

Tugba.yazgan@atauni.edu.tr, eilhan@ahievran.edu.tr, ercelik@atauni.edu.tr,
hbulut@firat.edu.tr

Abstract

In this study, some solitary wave solutions of the Wu-Zhang system are analyzed using the modified expansion function method and the sine-Gordon expansion method. Solitary wave solutions of this non-linear mathematical model consisting of hyperbolic and trigonometric function structures are get. Two and three dimensional, density graphics of the solitary solutions of the mathematical models are drawn by choosing the appropriate parameters. It was seen that all solution functions found as a result provide the mathematical model. In this study, Wolfram Mathematica software program was used for all mathematical calculations.

Keywords: The Sine-Gordon expansion method (SGEM), The modified expansion function method (MEFM), Wu-Zhang system.

1. Introduction

Each of the non-linear partial differential equations are mathematical models that can help to understand and solve problems such as physics, engineering, chemistry, biology. Recently, some approaches have been improved to search analytical solutions of several non-linear mathematical models. Some of those, the extended (G'/G) -expansion method [1], the Backlund transformation method [2], the simplified Hirota's method [3], the transformed rational function method [4], the modified simple equation method [5], the multiple exp-function method [6], the extended tanh method [7], the direct algebraic method [8], the Jacobi elliptic function method [9], the homogeneous balance method [10], the local fractional Riccati differential equation method [11], the improved Bernoulli sub-equation function method [12], Cornejo-Perez and Rosu method [13] among others. We used the sine-Gordon expansion

¹ Corresponding author

method [14] and the modified expansion function method [15]. for this purpose. Usually, diverse computational techniques have been improved to obtain solutions for different NLEEs [16-33].

In this study, it is planned to get the solitary solutions of Wu-Zhang system using MEFM [34] and the SGEM [35].

The Wu-Zhang system [36] is given as follows:

$$u_t + uu_x + v_x = 0, \quad (1)$$

$$v_t + (uv)_x + \frac{1}{3}u_{xxx} = 0. \quad (2)$$

In the equation system v the height of the water and u is the surface speed of water.

2. Methods

2.1. Overview of the MEFM

The general form of the non-linear partial differential equation (NPDE) is as follows:

$$P(u, u_x, u_t, u_{xxx}, v_x, v_t, uv_x, vu_x, K) = 0, \quad (3)$$

where $u = u(x, t)$ is unknown function, P is a polynomial in $u(x, t)$ and its derivatives.

Step 1: The travelling wave transformation is as follows:

$$u(x, t) = u(\eta), \quad \eta = r(x - ct). \quad (4)$$

If Eq. (4) is substituted in Eq. (3), the general form of non-linear ordinary differential equation (NODE) is obtained:

$$N(u, u^2, v, u'', K) = 0. \quad (5)$$

Step 2: We suppose the following wave equation to be the solution to Eq. (5):

$$u(\eta) = \frac{\sum_{j=0}^n A_j [e^{-g(\eta)}]^j}{\sum_{i=0}^m B_i [e^{-g(\eta)}]^i} = \frac{A_0 + A_1 e^{-g} + K + A_n e^{-ng}}{B_0 + B_1 e^{-g} + K + B_m e^{-mg}}, \quad (6)$$

where $A_j, B_i, (0 \leq j \leq n, 0 \leq i \leq m)$ are constants.

$$g'(\eta) = e^{-g(\eta)} + ke^{g(\eta)} + \lambda. \quad (7)$$

Eq.(7) has the following families of solutions [37]:

Family 1: When, $k \neq 0$ and $\lambda^2 - 4k > 0$,

$$\mathcal{G}(\eta) = \ln\left(\frac{-\sqrt{\lambda^2 - 4k}}{2k} \tanh\left(\frac{\sqrt{\lambda^2 - 4k}}{2}(\eta + E)\right) - \frac{\lambda}{2k}\right). \quad (8)$$

Family 2: When, $k \neq 0$ and $\lambda^2 - 4k < 0$,

$$\mathcal{G}(\eta) = \ln\left(\frac{\sqrt{-\lambda^2 + 4k}}{2k} \tan\left(\frac{\sqrt{-\lambda^2 + 4k}}{2}(\eta + E)\right) - \frac{\lambda}{2k}\right). \quad (9)$$

Family 3: When, $k = 0$, $\lambda \neq 0$ and $\lambda^2 - 4k > 0$,

$$\mathcal{G}(\eta) = -\ln\left(\frac{\lambda}{e^{\lambda(\eta+E)} - 1}\right). \quad (10)$$

Family 4: When, $k \neq 0$, $\lambda \neq 0$ and $\lambda^2 - 4k = 0$,

$$\mathcal{G}(\eta) = \ln\left(-\frac{2\lambda(\eta + E) + 4}{\lambda^2(\eta + E)}\right). \quad (11)$$

Family 5: When, $k = 0$, $\lambda = 0$ and $\lambda^2 - 4k = 0$,

$$\mathcal{G}(\eta) = \ln(\eta + E), \quad (12)$$

where $A_j, B_i, (0 \leq j \leq n, 0 \leq i \leq m)$, E, λ, k are coefficients and m, n positive integers got utilizing the balancing principle.

Step 3: Substituting Eq. (6) and its derivatives along with Eq. (7) into Eq. (5), an equation containing the polynomial is obtained. All coefficients are collected by collecting a series of algebraic equations of $e^{-\mathcal{G}(\eta)}$ have the same rank and make each sum equal to zero. To obtain new solutions for (3), the system of equations is solved with the help of Wolfram Mathematica program and the values of $A_j, B_i, (0 \leq j \leq n, 0 \leq i \leq m)$, E, λ, k coefficients are found. Considering the obtained coefficient values and equations (8-12), the solution functions that provide the equation (1) are obtained by replacing them in the equation (6).

2.2. Overview of the SGEM

Here, we give the analysis of the sine-Gordon equation [38]

$$u_{xx} - u_{tt} = v^2 \sin(u), \quad (13)$$

where $u = u(x, t)$ and $v \in \mathbb{R} - \{0\}$.

The travelling wave transformation $u = u(x, t) = u(\eta)$, $\eta = r(x - ct)$ on eq. (13), gives the following non-linear ordinary differential equation (NODE):

$$u'' = \frac{v^2}{r^2(1-c^2)} \sin(u), \quad (14)$$

where $u = u(\eta)$ and η stands for the width and k the velocity of the travelling wave respectively. Equation (14) can be simplified as follow forms:

$$\left[\left(\frac{u}{2} \right)' \right]^2 = \frac{v^2}{r^2(1-c^2)} \sin^2 \left(\frac{u}{2} \right) + q, \quad (15)$$

where q is the integration constant.

Substituting $q = 0$, $w(\eta) = \frac{u}{2}$ and $a^2 = \frac{v^2}{r^2(1-c^2)}$ into Eq. (15), yields

$$(w')^2 = a^2 \sin^2(w(\eta)), \quad (16)$$

substituting $a = 1$ into Eq. (16), yields

$$(w')^2 = \sin^2(w(\eta)), \quad (17)$$

From Eq. (17), we have the following four significant equations :

$$\sin(w(\eta)) = \sec h(\eta) \text{ or } \cos(w(\eta)) = -\tanh(\eta), \quad (18)$$

$$\sin(w(\eta)) = -i \operatorname{csch}(\eta) \text{ or } \cos(w(\eta)) = -\operatorname{coth}(\eta). \quad (19)$$

The solution of any non-linear partial differential equation (NPDE) is considered to be of the situations:

$$u(\eta) = \sum_{i=1}^n \tanh^{i-1}(\eta) [B_i \sec h(\eta) + A_i \tanh(\eta)] + A_0, \quad (20a)$$

$$u(\eta) = \sum_{i=1}^n \operatorname{coth}^{i-1}(\eta) [B_i \cot h(\eta) + iA_i \operatorname{csch}(\eta)] + A_0. \quad (20b)$$

According to Eq. (18) and (19), one way rewrite Eq. (20) as

$$u(w) = \sum_{i=1}^n \cos^{i-1}(w) [B_i \sin(w) + A_i \cos(w)] + A_0. \quad (21)$$

The value of n is defined by the balancing procedure of the highest order non-linear term and the highest order derivative. Substitution Eq. (21) and its possible derivatives into the NODE, gives an equation in different power of trigonometric functions “

$\sin^i(w) \cos^j(w)$, ($0 \leq i \leq n, 0 \leq j \leq n$)”. The coefficients of trigonometric functions of the similar sequence are summed and each sum is set equal to zero to obtain some algebraical equations. This set of algebraical equations is solved for the values of the corresponding

coefficients. Then the values of these coefficients are included in the equations. (20a) and (20b) to get the solutions of the given NPDE.

3. Applications

In this section, using two mathematical methods, we are obtain the solitary solutions of the Wu-Zhang system.

Let's think the following travelling wave transformation:

$$u = u(\eta), v = v(\eta), \eta = r(x - ct). \quad (22)$$

If the derivative terms required in equation (1) and (2) are obtained from the wave transformation and replaced.

Respectively,

$$-cr u' + r u u' + r v' = 0, \quad (23)$$

and

$$-cr v' + r v u' + r u v' + \frac{1}{3} r^3 u''' = 0. \quad (24)$$

Integrating Eq. (23), we get,

$$-cu + \frac{1}{2} u^2 + v = 0. \quad (25)$$

Simplifying Eq. (25), we have,

$$v' = cu' - uu'. \quad (26)$$

Substituting Eq. (26) into Eq. (24), we get,

$$2r^2 u'' + 9cu^2 - 3u^3 - 6c^2 u = 0. \quad (27)$$

3.1. Application of the MEFM

In this section, the MEFM is used to the Wu-Zhang system to obtain solitary solutions.

Balancing the highest power non-linear term and the highest derivative in Eq. (27), $n = m + 1$ gives the relationship.

Assume that $m = 1$. Then we have $n = 2$.

For m and n parameters, Eq. (6) is found as follows;

$$u(\zeta) = \frac{A_0 + A_1 e^{-\zeta} + A_2 e^{-2\zeta}}{B_0 + B_1 e^{-\zeta}}. \quad (28)$$

When the derivative term required in equation (27) is obtained from the expression (28), then polynomials equation of $e^{-\theta}$ is get. The algebraic equation system consisting of the coefficients of $e^{-\theta}$ is solved using mathematica program to get the following conditions:

Case-1:

$$A_0 = cB_0 - \frac{\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2}}{\lambda^2 - 4k},$$

$$A_1 = \frac{-\lambda\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2}B_1 + B_0\left(-2\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2} + c\lambda(\lambda^2 - 4k)B_1\right)}{\lambda(\lambda^2 - 4k)B_0},$$

$$A_2 = -\frac{2c^2\lambda B_0 B_1}{\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2}}, s = \frac{\sqrt{3}c}{\sqrt{\lambda^2 - 4k}}.$$

The following solutions are obtained after these coefficients are put in Eq. (28),

Family 1: When $k \neq 0$, $\lambda^2 - 4k > 0$, solution of equation (1),

$$u_1(x, t) = \frac{c\left(\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2}(\lambda + \tau) - c\lambda B_0(\lambda^2 - 4k + \lambda\tau)\right)}{\sqrt{c^2\lambda^2(\lambda^2 - 4k)B_0^2}(\lambda + \tau)}, \quad (29)$$

$$v_1(x, t) = \frac{(4c^2k)}{\left(\left(1 + \text{Cosh}\left[\sqrt{3}c\eta + E\sqrt{\lambda^2 - 4k}\right]\right)(\lambda + \tau)\right)^2}.$$

Where $\tau = \left(\sqrt{\lambda^2 - 4k}\text{Tanh}\left[\frac{1}{2}\sqrt{3}c\eta\left(E + \sqrt{\lambda^2 - 4k}\right)\right]\right)$.




Figure-1. The three dimensional, density graphic and two dimensional graphic surfaces of Eq. (29) in $k=1, \lambda=3, E=0.75, c=1, B_0=0.35$ and $t=1$.

Family 2: When $k \neq 0, \lambda^2 - 4k < 0$,

$$u_2(x,t) = \frac{c}{c\lambda B_0 \kappa} (\lambda + \kappa \gamma)^{-1} (c\lambda B_0 \kappa (\lambda + \kappa \gamma) - c\lambda B_0 (\lambda^2 - 4k + \lambda \kappa \gamma)), \quad (30)$$

$$v_2(x,t) = 4k c^2 \left((1 + \text{Cosh}[\sqrt{3} c \eta + E \kappa]) \left(\lambda + \kappa \text{Tanh} \left[\frac{1}{2} (\sqrt{3} c \eta + E \kappa) \right] \right) \right)^{-2},$$

where $\left(\kappa = \sqrt{\lambda^2 - 4k}, \gamma = \text{Tanh} \left[\frac{\kappa}{2} (E + \eta) \right] \right)$.




Figure-2. The three dimensional, density and two dimensional graphic surfaces of Eq. (30) in $k = 1, \lambda = 3, c = 1, B_0 = 0.35, E = 0.75$ and $t = 1$.

Family 3: When $k = 0, \lambda \neq 0$ and $\lambda^2 - 4k > 0$,

$$u_3(x, t) = c - \frac{\text{Coth} \left[\frac{1}{2} \lambda \left(E + \frac{\sqrt{3} c \eta}{\sqrt{\lambda^2}} \right) \right] \sqrt{c^2 \lambda^4 B_0^2}}{\lambda^2 B_0}, \quad (31)$$

$$v_3(x, t) = - \frac{c^2}{-1 + \text{Cosh} \left[\lambda \left(E + \frac{\sqrt{3} c \eta}{\sqrt{\lambda^2}} \right) \right]}.$$




Figure-3. The three dimensional, density and two dimensional graphic surfaces of Eq. (31) in $k=1$, $\lambda=3$, $c=1$, $E=0.75$ $B_0=0.35$ and $t=1$.

According to the cases Family 4 and Family 5, the solution can not be get. Because, the solution function u is calculated as undefined due to the $\lambda^2 - 4k = 0$ term.

Case-2:

$$A_0 = -\sqrt{\frac{2}{3}} \sqrt{s^2(\lambda^2 - 2k)B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4}},$$

$$A_1 = \frac{\sqrt{s^2(\lambda^2 - 2k)B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4}} \left(\sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4} - s^2\lambda B_0(\lambda B_0 + 2kB_1) \right)}{\sqrt{6}s^2\lambda kB_0^2},$$

$$A_2 = \frac{\left(-s^2\lambda^2 B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4} \right) \sqrt{s^2(\lambda^2 - 2k)B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4}}}{\sqrt{6}s^2\lambda kB_0^3},$$

$$v_5(x,t) = \frac{\left(2s^2(\lambda^2 - 4k)k \operatorname{Sech} \left[\frac{1}{2} \sqrt{\lambda^2 - 4k} \left(E + sx + \frac{\tau}{2\sqrt{6}skB_0^3} \right) \right] \right)^2}{\left(3 \left(\lambda + \sqrt{\lambda^2 - 4k} \operatorname{Tanh} \left[\frac{1}{2} \sqrt{\lambda^2 - 4k} \left(E + sx + \frac{1}{(2\sqrt{6}skB_0^3)\tau} \right) \right] \right) \right)^2},$$

where, $\tau = \left(t \left(-s^2(\lambda^2 - 4k)B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4} \right) \sqrt{s^2(\lambda^2 - 2k)B_0^2 + \sqrt{s^4\lambda^2(\lambda^2 - 4k)B_0^4}} \right)$.




Figure-5. The three dimensional, density and two dimensional graphic surfaces of Eq. (33) in $k = 1, c = 1, \lambda = 3, s = 1, B_0 = 0.35, E = 0.75$ and $t = 1$.




Figure-6. The three dimensional, density and two dimensional graphic surfaces of Eq. (34) in $k = 1, c = 1, \lambda = 3, s = 1, B_0 = 0.35, E = 0.75$ and $t = 1$.

3.2. Application of the SGEM

In this section, solitary solutions are obtained by applying the SGEM method to the Wu-Zhang system.

Balancing the highest power non-linear term and the highest derivative in Eq. (26), gives $n = 1$

Equation (21) is written as follows according to n value,

$$u(w) = B_1 \sin(w) + A_1 \cos(w) + A_0. \quad (35)$$

Substituting Eq. (35) and its second derivative along with Eq. (17) into Eq. (27), gives an equation in power of trigonometric functions. We collect a set of algebraical equations by equating the summations of the coefficients of the trigonometric functions with the same power to zero. The set of algebraical equations is simplified to obtain the values of the parameters included. The values of parameters are then substituted into Eq. (20a) and (20b) with fixed value of m to get the solutions of Eq. (1).

Case-1: When,

$$A_0 = -\frac{k}{\sqrt{3}}, A_1 = -\frac{k}{\sqrt{3}}, B_1 = \frac{ik}{\sqrt{3}}, c = -\frac{k}{\sqrt{3}},$$

we have the following compound non-topological and topological kink-type soliton:



$$u_7(x,t) = \frac{k \left(-1 + i \operatorname{Sech} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] + \operatorname{Tanh} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] \right)}{\sqrt{3}}, \quad (36)$$

$$v_7(x,t) = -\frac{k^2}{3 + 3i \operatorname{Sinh} \left[k \left(x + \frac{tk}{\sqrt{3}} \right) \right]},$$

and the singular soliton,

$$u_8(x,t) = -\frac{k \left(-1 + \operatorname{Coth} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] + \operatorname{Csch} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] \right)}{\sqrt{3}}, \quad (37)$$

$$v_8(x,t) = -\frac{1}{6} k^2 \operatorname{Csch} \left[\frac{1}{2} k \left(x + \frac{tk}{\sqrt{3}} \right) \right]^2.$$






Figure-7. The three dimensional, density and two dimensional graphic surfaces of the imaginary and real part of the Eq.(36) respectively in $k = 1$, $t = 1$.




Figure-8. The three dimensional, density and two dimensional graphic surfaces of the equation (37) in $k = 1$, $t = 1$.

Case-2: When,

$$A_0 = \frac{2k}{\sqrt{3}}, A_1 = -\frac{2k}{\sqrt{3}}, B_1 = 0, c = \frac{2k}{\sqrt{3}},$$

we get the following topological kink-type soliton:

$$u_9(x,t) = \frac{2k}{\sqrt{3}} \left(1 + \text{Tanh} \left[k \left(x - \frac{2tu}{\sqrt{3}} \right) \right] \right),$$

$$v_9(x,t) = \frac{2}{3} k^2 \text{Sech} \left[k \left(x - \frac{2tk}{\sqrt{3}} \right) \right]^2.$$
(38)




Figure-9. The three dimensional, density and two dimensional graphic surfaces of Eq.(38) in $k=1$, $t=1$.

4. Results and Discussion

The modified exp function method and the Sine-Gordon expansion method have been successfully employed to secure the wave solutions of an important non-linear model; the Wu-Zhang system. Various wave solutions obtained by using these powerful schemes have been reported in this study. On the other hand, when we compare the results obtained by using two methods in this article with the results obtained in [39], some new travelling wave solutions have been presented to the literature. The hyperbolic function solutions acquired by using the

modified expansion function method of Wu-Zhang system are obtained from the following coefficients.

$$A_0 = cB_0 - \frac{\sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2}{\lambda^2 - 4k},$$

$$A_1 = \frac{-\lambda \sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2 B_1 + B_0 \left(-2\sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2 + c\lambda (\lambda^2 - 4k) B_1 \right)}{\lambda (\lambda^2 - 4k) B_0},$$

$$A_2 = -\frac{2c^2 \lambda B_0 B_1}{\sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2}, s = \frac{\sqrt{3}c}{\sqrt{\lambda^2 - 4k}}.$$

From these coefficients, diverse solution functions are obtained according to the family-1-2-3 conditions. For example;

$$u_1(x, t) = \frac{c \left(\frac{\sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2 \left(\lambda + \sqrt{\lambda^2 - 4k} \operatorname{Tanh} \left[\frac{1}{2} \sqrt{3} c \eta (E + \sqrt{\lambda^2 - 4k}) \right] \right) - c\lambda B_0 \left(\lambda^2 - 4k + \lambda \sqrt{\lambda^2 - 4k} \operatorname{Tanh} \left[\frac{1}{2} \sqrt{3} c \eta (E + \sqrt{\lambda^2 - 4k}) \right] \right) \right)}{\sqrt{c^2 \lambda^2 (\lambda^2 - 4k)} B_0^2 \left(\lambda + \sqrt{\lambda^2 - 4k} \operatorname{Tanh} \left[\frac{1}{2} \sqrt{3} c \eta (E + \sqrt{\lambda^2 - 4k}) \right] \right)},$$

$$v_1(x, t) = \frac{(4c^2 k)}{\left(\left(1 + \operatorname{Cosh} \left[\sqrt{3} c \eta + E \sqrt{\lambda^2 - 4k} \right] \right) \right)^2 \left(\left(\lambda + \sqrt{\lambda^2 - 4k} \operatorname{Tanh} \left[\frac{1}{2} (\sqrt{3} c \eta + E \sqrt{\lambda^2 - 4k}) \right] \right) \right)}.$$

Some of the solutions of the Wu-Zhang system according to the coefficients found according to the other method, sine-Gordon expansion method, are as follows.

$$A_0 = -\frac{k}{\sqrt{3}}, A_1 = -\frac{k}{\sqrt{3}}, B_1 = \frac{ik}{\sqrt{3}}, c = -\frac{k}{\sqrt{3}},$$

$$u_7(x, t) = \frac{k \left(-1 + i \operatorname{Sech} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] + \operatorname{Tanh} \left[k \left(x + \frac{tu}{\sqrt{3}} \right) \right] \right)}{\sqrt{3}},$$

$$v_7(x,t) = -\frac{k^2}{3 + 3i \operatorname{Sinh} \left[k \left(x + \frac{tk}{\sqrt{3}} \right) \right]}.$$

The results acquired successfully in our study are thought to have an important physical meaning for the dynamical system. Periodic features are needed to make an estimate of the prospective behavior of the system.

5. Conclusions

In this article, we construct various wave solutions to the Wu-Zhang system by using the modified expansion function method and the sine-Gordon expansion method. We successfully get topological kink-type, non-topological, singular soliton and trigonometric function solutions. When the graphics of the obtained solutions of the non-linear differential equation are analyzed, they physically inform the motion pattern of the wave. When the graphs of the solution functions obtained as a result of the methods applied to the equations are physically interpreted, it is observed that the movements intensify according to the characteristic feature of the solution function as time progresses and pigs in certain points. The graphics indicate that the solution functions obtained have periodic features. It is benefit to obtain functions with such features. Because it is quite easy to physically interpret equations with periodic function properties. It helps us to easily comment on the motion model within the desired range.

The reported results show that the two methods are very efficient and suitable mathematical tools for solving non-linear partial differential equations.

References

- [1] S. Kumar, K. Singh, and R. K. Gupta, [Coupled Higgs field equation and Hamiltonian amplitude equation: Lie classical approach and \(G'/G\)-expansion method](#), *Pramana*, **79** (2012), 41-60.
- [2] D. C. Lu, B. J. Hong, and L. Tian, [Backlund transformation and n-soliton-like solutions to the combined KdV-Burgers equation with variable coefficients](#), *International Journal of Nonlinear Science*, **2** (2006), 3-10.
- [3] M. T. Darvishi, and M. Najafi, [Some exact solutions of the \(2+ 1\)-dimensional breaking soliton equation using the three-wave method](#), *International Journal of Computational and Mathematical Sciences*, **6** (2012), 13-16.

- [4] R. Seadawy, [Travelling-wave solutions of a weakly non-linear two-dimensional higher-order Kadomtsev-Petviashvili dynamical equation for dispersive shallow-water waves](#), The European Physical Journal Plus, **132** (2017), 29.
- [5] A. J. A. M. Jawad, M. D. Petković, and A. Biswas, [Modified simple equation method for non-linear evolution equations](#), Applied Mathematics and Computation, **217** (2010), 869-877.
- [6] W. X. Ma, T. Huang, and Y. Zhang, [A multiple exp-function method for non-linear differential equations and its application](#), Physica Scripta, **82** (2010), 065003.
- [7] M. A. Abdou, [The extended tanh method and its applications for solving non-linear physical models](#), Applied mathematics and computation, **190** (2007), 988-996.
- [8] H. Willy, et al., [Exact solitary wave solutions of non-linear evolution and wave equations using a direct algebraic method](#), Journal of Physics A: Mathematical and General, **19** (1986), 607.
- [9] L. Shikuo, et al., Jacobi elliptic function expansion method and periodic wave solutions of non-linear wave equations, Phys Lett A, **289** (2001), 69-74.
- [10] M. Wang, Z. Yubin, and L. Zhibin, [Application of a homogeneous balance method to exact solutions of non-linear equations in mathematical physics](#), Physics Letters A, **216** (1996), 67-75.
- [11] H. M. Baskonus, and H. Bulut, [Exponential prototype structures for \(2+ 1\) dimensional Boiti-Leon-Pempinelli systems in mathematical physics](#), Waves in Random and Complex Media, **26** (2016), 189-196.
- [12] H. M. Baskonus, and H. Bulut, [On the complex structures of Kundu-Eckhaus equation via improved Bernoulli sub-equation function method](#), Waves in Random and Complex Media, **25** (2015), 720-728.
- [13] O. Cornejo-Pérez, and H. C. Rosu, [Non-linear second order Ode's: factorizations and particular solutions](#), Progress of theoretical physics, **114** (2005), 533-538.
- [14] Y. Chen, and Y. Zhenya, [New exact solutions of \(2+ 1\)-dimensional Gardner equation via the new sine-Gordon equation expansion method](#), Chaos, Solitons & Fractals, **26** (2005), 399-406.
- [15] J. H. He, XH Wu Chaos, Solitons, Fractals, **30** (2006), 700.
- [16] F. Gao, X. J. Yang, and Y. F. Zhang, [Exact traveling wave solutions for a new non-linear heat transfer equation](#), Thermal Science, **21** (2017), 1833-1838.
- [17] H. Bulut, T. A. Sulaiman, and B. Demirdag, [Dynamics of soliton solutions in the chiral non-linear Schrödinger equations](#), Non-linear Dynamics, **91** (2018), 1985-1991.

- [18] T. A. Sulaiman, et al, [Investigation of various soliton solutions to the Heisenberg ferromagnetic spin chain equation](#), Journal of Electromagnetic Waves and Applications, **32** (2018), 1093-1105.
- [19] Z. Y. Zhang, [Jacobi elliptic function expansion method for the modified Korteweg-de Vries-Zakharov-Kuznetsov and the Hirota equations](#), Romanian Journal of Physics, **60** (2015), 1384-1394.
- [20] C. Cattani, et al., [On the soliton solutions to the Nizhnik-Novikov-Veselov and the Drinfel'd-Sokolov systems](#), Optical and Quantum Electronics, **50** (2018) 138.
- [21] H. Bulut, T. Akturk, and Y. Gurefe, [Traveling wave solutions of the \(N+ 1\)-dimensional sine-cosine-Gordon equation](#), AIP Conference Proceedings, **1637** American Institute of Physics, 2014.
- [22] A. Yokus, et al., [Numerical simulation and solutions of the two-component second order KdV evolutionary system](#), Numerical Methods for Partial Differential Equations, **34** (2018), 211-227.
- [23] M. Inc, Y. Abdullahi, A. I. Aliyu, and D. Baleanu, [Soliton solutions and stability analysis for some conformable non-linear partial differential equations in mathematical physics](#), Optical and Quantum Electronics, **50** (2018), 190.
- [24] H. M. Baskonus, H. Bulut, and T. A. Sulaiman, [Investigation of various travelling wave solutions to the extended \(2+ 1\)-dimensional quantum ZK equation](#), The European Physical Journal Plus, **132** (2017), 1-8.
- [25] E. M. E. Zayed, and S. H. Ibrahim, [Exact solutions of non-linear evolution equations in mathematical physics using the modified simple equation method](#), Chinese Physics Letters, **29** (2012), 060201.
- [26] H. Bulut, T. A. Sulaiman, and H. M. Baskonus, [On the new soliton and optical wave structures to some non-linear evolution equations](#), The European Physical Journal Plus, **132** (2017), 459.
- [27] K. Hosseini, et al., [Travelling wave solutions of the Korteweg-de Vries equation with dual-power law non-linearity using the improved tan \(\$\phi\$ \(\$\xi\$ \)/2\)-expansion method](#), Optik, **156** (2018), 498-504.
- [28] H. Bulut, et al., [Novel hyperbolic behaviors to some important models arising in quantum science](#), Optical and Quantum Electronics, **49** (2017), 349.
- [29] H. M. Baskonus, T. A. Sulaiman, and H. Bulut, [On the novel wave behaviors to the coupled non-linear Maccari's system with complex structure](#), Optik, **131** (2017), 1036-1043.

- [30] M. G. Hafez, M. N. Alam, and M. A. Akbar, Application of the $\exp(-\Phi(\eta))$ expansion method to find exact solutions for the solitary wave equation in an unmagnetized dusty plasma, *World Applied Sciences Journal*, **32** (2014), 2150-2155.
- [31] H. Bulut, et al., [New solitary and optical wave structures to the \(1+ 1\)-dimensional combined KdV–mKdV equation](#), *Optik*, **135** (2017), 327-336.
- [32] K. Khan, and M. A. Akbar, [Application of exp-expansion method to find the exact solutions of modified Benjamin-Bona-Mahony equation](#), *World Applied Sciences Journal*, **24** (2013), 1373-1377.
- [33] H. M. Baskonus, T. A. Sulaiman, and H. Bulut, [New solitary wave solutions to the \(2+1\)-dimensional Calogero–Bogoyavlenskii–Schiff and the Kadomtsev–Petviashvili hierarchy equations](#), *Indian Journal of Physics*, **91** (2017), 1237-1243.
- [34] H. M. Baskonus, H. Bulut, and A., [On the complex and hyperbolic structures of the longitudinal wave equation in a magneto-electro-elastic circular rod](#), *Smart Materials and Structures*, **25** (2016), 035022.
- [35] Y. Chen, and Z. Yan, [New exact solutions of \(2+ 1\)-dimensional Gardner equation via the new sine-Gordon equation expansion method](#), *Chaos, Solitons & Fractals*, **26** (2005), 399-406.
- [36] H. Jafari, et al, [On the exact solutions of non-linear long-short wave resonance equations](#), *Rom. Rep. Phys.*, **67** (2015), 762-772.
- [37] H. Naher, and F. A. Abdullah, [New generalized and improved \(G'/G\)-expansion method for non-linear evolution equations in mathematical physics](#), *Journal of the Egyptian Mathematical Society*, **22** (2014), 390-395.
- [38] M. A. Abdelrahman, E. H. Zahran, and M. M. Khater, [The Exp \$\(-\phi\(\xi\)\)\$ -expansion method and its application for solving non-linear evolution equations](#), *International Journal of Modern Nonlinear Theory and Application*, **4** (2015), 37.
- [39] M. Eslami, and H. Rezazadeh, [The first integral method for Wu–Zhang system with conformable time-fractional derivative](#), *Calcolo*, **53** (2016), 475-485.